



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 333/38, A61K 31/38, 31/47, 31/55, C07D 307/68, 409/12		A1	(11) International Publication Number: WO 97/06158
			(43) International Publication Date: 20 February 1997 (20.02.97)
(21) International Application Number: PCT/GB95/01896 (22) International Filing Date: 10 August 1995 (10.08.95) (71) Applicant (for all designated States except MN): ASTRA AB [SE/SE]; S-151 85 Södertälje (SE). (71) Applicant (for MN only): ASTRA PHARMACEUTICALS LIMITED [GB/GB]; Home Park, Kings Langley, Hertfordshire, WD4 8DH (GB). (72) Inventors: MacDONALD, James, Edwin; 92 Canfield Road, Pittsford, NY 14534 (US). SHAKESPEARE, William, Calvin; 127 Raleigh Street, Rochester, NY 14620 (US). MURRAY, Robert, John; 172 Lac Kine Drive, Brighton, NY 14618 (US). MATZ, James, Russell; 60 Waterford Way, Fairport, NY 14450 (US). (74) Agent: TEUTEN, Andrew, John; Astra Patents, Loughborough, P.O. Box 86, Loughborough, Leicestershire LE11 0RJ (GB).		(81) Designated States: AM, AT, BB, BG, BR, BY, CH, CN, CZ, DE, DK, EE, ES, GB, GE, HU, IS, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, UZ, VN, ARIPO patent (KE, MW, SD, SZ, UG), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: BICYCLIC AMIDINE DERIVATIVES USEFUL IN THERAPY			
(57) Abstract			
<p>There are provided compounds of formula (I), wherein D represents a five membered heterocyclic aromatic ring containing 1 to 4 heteroatoms selected from O, N or S, optionally substituted at a carbon atom by halogen, trifluoromethyl, alkyl C1 to 6, nitro or cyano, and which is connected to the remainder of the compound of formula (I) through a carbon atom; A represents N(X) or CH(-(CH₂)_m-NXY); U represents NH, O or CH₂; V represents (CH₂)_a; W represents (CH₂)_b; a, b, m, X and Y are as defined in the specification, together with processes for their preparation and compositions containing them. Compounds of formula (I) are nitric oxide synthetase inhibitors and are useful in therapy.</p>		<p style="text-align: right;">(I)</p>	

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Bicyclic amidine derivatives useful in therapy

5 This invention relates to bicyclic amidine derivatives, processes for their preparation, compositions containing them and their use in therapy.

Certain amidine derivatives have been described for use in therapeutic applications. N-Phenyl amidine derivatives have been described for use in the treatment of diabetes in US Patent No. 3669974 (USV Pharmaceutical Corp.) and
10 UK Patent Application 2226562 (Boots). N'N''disubstituted amidines are described for use in the treatment of hypertension, depression and hallucinogenic states in International Patent Application WO 92/04054 (University of Oregon). The use of certain amidines and symmetric bisamidines as analgesics, in the treatment of inflammation and in the treatment of hypertension is described in Belgian Patent
15 No. 717740 and UK Patent No. 1180629 (both of Delalande). Amidine derivatives have also been described for use as herbicides in German Patent Application DE-OS-2321330 (Bayer).

The use of inhibitors of nitric oxide synthetase in the treatment of disease has
20 also been described, for example, in International Patent Applications WO 94/12163 (Abbott), WO 93/13066 and WO 94/12165 (both of Wellcome) and European Patent Applications 446699 (Merrell Dow), 547558 and 558468 (both of Washington University). The use of nitric oxide synthetase inhibitors in therapy is also described in WO 95/00505, WO 95/09619, WO 95/09621 (all of Wellcome), WO
25 95/10266 (Otsuka), WO 95/11231 and WO 95/11014 (both of Searle).


The applicant has previously described the use of guanidine derivatives and amidine derivatives which are inhibitors of nitric oxide synthetase in the treatment
inter alia of neurodegenerative disease (WO 94/21621, WO 95/05363).

30

-2-

We have now found a new group of bicyclic amidine derivatives that possesses useful pharmaceutical activity.

According to a first aspect of the invention, we provide a compound of formula
5 I



10

wherein

D represents a five membered heterocyclic aromatic ring containing 1 to 4 heteroatoms selected from O, N or S, optionally substituted at a carbon atom by halogen, trifluoromethyl, alkyl C1 to 6, nitro or cyano, and which is connected to
15 the remainder of the compound of formula I through a carbon atom;

A represents N(X) or CH(-(CH₂)_m-NXY);

U represents NH, O or CH₂;

V represents (CH₂)_a;

W represents (CH₂)_b;

20 a and b independently represent an integer 0 to 3,

provided that a+b is in the range 1 to 3;

X and Y independently represent hydrogen, alkyl C1 to 6, or the group -(CH₂)_nQ, or -NXY represents piperidinyl, pyrrolidinyl, morpholinyl or tetrahydroisoquinolinyl;

Q represents biphenyl or phenyl optionally substituted by one or more groups
25 selected from alkyl C1 to 6, alkoxy C1 to 6, perfluoroalkyl C1 to 6, halogen, nitro or cyano;

m represents an integer 0 to 5;

n represents an integer 0 to 6;

or the chain U-V-A-W is as defined above save that it may be unsaturated,


30 or the chain U-V-A-W may represent -NH-CH₂-CH₂-O- substituted at a carbon atom by the group -(CH₂)_m-NXY, wherein m, X and Y are as defined above,

-3-

and pharmaceutically acceptable salts thereof.

A preferred group of compounds of formula I is defined by formula IA:

5



wherein

- T represents a C_{3-5} saturated or unsaturated alkylene chain substituted by $-(CH_2)_m-$
 10 NXY; $-O-(CH_2)_2-NH-$ substituted by $-(CH_2)_m-NXY$; or $-U-(CH_2)_a-N(X)-(CH_2)_b-$;
 X and Y independently represent hydrogen, alkyl C1 to 6, or the group $-(CH_2)_nQ$,
 or -NXY represents piperidinyl, pyrrolidinyl, morpholinyl or tetrahydroisoquinolinyl;
 Q represents phenyl optionally substituted by alkyl C1 to 6, alkoxy C1 to 6,
 trifluoromethyl, halogen, nitro or cyano;
 15 and U, m, n, a, b and D are as defined above,
 save that when T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$ and X represents $-(CH_2)_nQ$, n
 represents an integer 0 to 5,
 and pharmaceutically acceptable salts thereof.

- 20 We prefer that D represents a five-membered heterocyclic aromatic ring
 containing one heteroatom selected from O, N or S, optionally substituted at a
 carbon atom by halogen. We particularly prefer that D represents thienyl, furyl or
 pyrrolyl, especially thienyl or furyl, more especially thienyl and most especially 2-
 thienyl.

25

We prefer that T represents a C_{3-5} saturated or unsaturated alkylene chain
 substituted by $-(CH_2)_m-NXY$, particularly a C_{3-5} saturated alkylene chain substituted
 by $-(CH_2)_m-NXY$, especially a C_{3-4} saturated alkylene chain substituted by $-(CH_2)_m-$
 NXY.

30

-4-

When T represents a $C_{3,5}$ saturated or unsaturated alkylene chain substituted by $-(CH_2)_m-NXY$; or $-O-(CH_2)_2-NH-$ substituted by $-(CH_2)_m-NXY$, we prefer that X and Y independently represent hydrogen, alkyl C1 to 6 or the group $-(CH_2)_nQ$. We particularly prefer that X and Y independently represent hydrogen, methyl, ethyl or
 5 the group $-(CH_2)_nQ$ and especially that one of X and Y represents hydrogen and the other represents hydrogen or the group $-(CH_2)_nQ$.

We prefer that m represents 0 or 1, especially 0.

10 When T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$, we prefer U to represent CH_2 .

When T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$, we prefer that $a+b$ is 1 or 2.


When T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$, we prefer that X represents
 15 hydrogen, alkyl C1 to 6 or the group $-(CH_2)_nQ$.

When X and/or Y represent $-(CH_2)_nQ$, we prefer that n represents 0, 1 or 2, especially 1.

20 We prefer that Q represents phenyl optionally substituted by alkyl C1 to 6 or halogen, although we particularly prefer that Q represents unsubstituted phenyl.

According to the invention, we further provide a process for the preparation of compounds of formula I, and pharmaceutically acceptable salts thereof, which
 25 comprises:

(a) preparing a compound of formula I by reacting a corresponding compound of formula II




30

wherein U, V, A and W are as defined above,


-5-

with a compound of formula III



5 wherein D is as defined above and L is a leaving group;

(b) preparing a compound of formula I by reacting a corresponding compound of formula IV




wherein U, V, A and W are as defined above and HA is an acid,
with a compound of formula V



wherein D is as defined above;

(c) preparing a compound of formula I in which A represents N(X) and X represents alkyl C1 to 6 or the group $-(\text{CH}_2)_n\text{Q}$ by reacting a corresponding compound of formula I in which X represents hydrogen with a compound of

20 formula VI




wherein R^9 represents alkyl C1 to 6 or the group $-(\text{CH}_2)_n\text{Q}$ and L is a leaving group;

25 (d) preparing a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m\text{---} \text{NXY})$ and at least one of X and Y represents alkyl C1 to 6 or the group $-(\text{CH}_2)_n\text{Q}$ by reacting a corresponding compound of formula I in which one or both of X and Y represents hydrogen with a compound of formula VI;


(e) preparing a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m\text{---} \text{NXY})$ and m represents an integer 1 to 5, by reduction of a corresponding
30 compound of formula VII

-6-




5 wherein U, V, W, X, Y and D are as defined above;

(f) preparing of a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$ and both X and Y represent hydrogen, by reduction of a corresponding compound of formula VIII




wherein U, V, W, m and D are as defined above;

(g) preparing a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, X represents hydrogen and m represents an integer 1 to 5, by reduction of a corresponding compound of formula IX



wherein U, V, W, D and Y are as defined above;


(h) preparing a compound of formula I wherein A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, one of X and Y represents hydrogen, and the other represents $-(\text{CH}_2)_n\text{Q}$ in which n represents an integer 1 to 6, by reduction of a corresponding compound of formula X



30. wherein Q, m, U, V, W and D are as defined above;


-7-

- (i) preparing a compound of formula I wherein A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, one of X and Y represents hydrogen, and the other represents $-(\text{CH}_2)_n\text{Q}$ in which n represents an integer 1 to 6, by reduction of a compound of formula XI



wherein Q, m, U, V, W and D are as defined above; or

- (j) preparing a compound of formula I in which A represents $\text{CH}(-\text{NXY})$ and
 10 X represents hydrogen by reduction of a corresponding compound of formula XII



- 15 wherein U, V, W, D and Y are as defined above;

and where desired or necessary converting the resultant compound of formula I, or another salt thereof, to a pharmaceutically acceptable salt thereof, or vice versa.

- 20 In process (a), the reaction will take place on stirring a mixture of the reactants in a suitable solvent, for example a lower alkanol e.g. ethanol, isopropanol or tertiary butanol, at a temperature between room temperature and the reflux temperature of the solvent. The reaction time will depend inter alia on the solvent and the nature of the leaving group, and may be up to 48 hours, however it will
 25 typically be from 1 to 5 hours. Suitable leaving groups that L may represent include thioalkyl, sulphonyl, trifluorocarbon sulphonyl, halide, alkyl and aryl alcohols and tosyl groups; others are recited in 'Advanced Organic Chemistry', J. March (1985) 3rd Edition, McGraw-Hill on page 315 and are well known in the art.

- 30 In process (b), the reaction is preferably performed by refluxing a mixture of the two compounds for several hours in the presence of a suitable solvent whereby the

-8-

reaction temperature is high enough so that condensation takes place readily, but not sufficiently high to decompose the amidine formed. The reaction temperature can vary from room temperature to about 250 °C, although it is preferable to perform the reaction at temperatures from about 100 °C to 200 °C. We find that o-
5 dichlorobenzene is a particularly suitable solvent and it is useful to add 4-dimethylaminopyridine as a catalyst. On cooling, two layers form, the solvent may be decanted, and the reaction worked up by addition of aqueous base. Alternatively, where the reactants are soluble in the solvent, the solvent may be evaporated off under vacuum and the reaction mixture worked up by addition of
10 water. The acid HA may be an organic or inorganic acid, for instance, hydrochloric, hydrobromic, hydroiodic, sulphuric, nitric, phosphoric, acetic, lactic, succinic, fumaric, malic, maleic, tartaric, citric, benzoic or methanesulphonic acid.

In process (c), the reaction will take place under standard conditions, for
15 example by reacting the two compounds in an inert solvent under basic conditions at room temperature for a period of up to 12 hours. We have frequently found it desirable to treat the amine with NaH before reacting with the compound of formula VI. We prefer that L represents halide, particularly bromide.

20 Process (d) may be performed under conditions analogous to those described above for process (c).

In process (e), the reduction may be performed by treatment with diborane in an inert solvent e.g. THF. Alternative although less preferred reagents which may be
25 suitable include lithium aluminium hydride and reagents for catalytic hydrogenation e.g. H₂ on Pd/C. Further details of the reaction conditions for use of these reactions may be obtained by reference to J. March "Advanced Organic Chemistry" on page 1099, including the references cited therein.

30 In process (f), the reduction reaction may be performed under a number of conditions, for example those described in J March "Advanced Organic Chemistry"

-9-

on pages 1103-1104. These include catalytic hydrogenation, use of Zn, Sn or Fe metal, $\text{AlH}_3\text{-AlCl}_3$, sulphides and others. We prefer to perform the reaction by hydrogenation at atmospheric pressure for 3-6 hours in the presence of a palladium and carbon catalyst.

5


In processes (g), (i) and (j), the reduction may be performed by treating the compound with sodium borohydride or sodium cyanoborohydride under standard conditions.

10 In process (h) the reaction may be performed under conditions analogous to those described above for process (e).

Salts of compounds of formula I may be formed by reacting the free base or a salt, enantiomer, tautomer or protected derivative thereof, with one or more
 15 equivalents of the appropriate acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, eg water, dioxan, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed in vacuo or by freeze drying. The reaction may be a metathetical process or it may be carried out on an ion exchange resin.

20

The compounds of formula II may be prepared by reduction of a corresponding compound of formula XIII



X | | |

25

wherein U, V, A and W are as defined above.

The reduction reaction may be performed under analogous conditions to those described above for process (f).

30

-10-

Certain compounds of formula II are either known or may be prepared by conventional methods known per se. Other compounds of formula II may be prepared from known compounds with simpler substituent groups by following analogous processes to those described above for processes (c) to (j). For example, by analogy with process (j) above, we find it convenient to prepare certain compounds of formula XIII in which A represents CH(-NXY) and X represents hydrogen by reduction of the corresponding imine formed by reaction of a compound of formula NH₂Y with the nitrated bicyclic ketone.

Compounds of formula IV may be prepared by analogous processes to those described for the preparation of compounds of formula II. Compounds of formula IV may be converted to corresponding compounds of formula II by treatment with a base. Compounds of formula II may be converted to corresponding compounds of formula IV by treatment with a protic acid HA, for example one of those listed above.

Compounds of formula III are either known or may be prepared by known methods. For example, compounds of formula III in which L represents thioalkyl may be prepared by treatment of the corresponding thiamide of formula XIV



wherein D is as defined above,
with an alkyl iodide.

Compounds of formula VII, VIII, IX, X, XI and XII may be prepared by analogous processes to those described for the preparation of compounds of formula I. Such compounds may be readily prepared from compounds with simpler substituent groups by conventional methods e.g. formation of an amide (VII, X) by reaction of an amine with a carboxylic acid or activated derivative thereof or formation of an imine (IX, XI, XII) by reaction of an amine with an aldehyde.

-11-

Compounds of formula V, VI, XIII and XIV are either known or may be prepared by conventional methods known per se.

It will be apparent to a person skilled in the art that it may be desirable to
5 protect an amine or other reactive group using a protecting group as described in
the standard text "Protecting groups in Organic Synthesis", 2nd Edition (1991) by
Greene and Wuts. Amine-protecting groups which may be mentioned include
alkyloxycarbonyl C2 to 7, eg *t*-butoxycarbonyl, phenylalkyloxycarbonyl C8 to 13, eg
benzyloxycarbonyl or preferably trifluoroacetate. Deprotection will normally take
10 place on treatment with aqueous base, acid or by treatment with hydrogen.

The compounds of the invention and intermediates may be isolated from their
reaction mixtures by standard techniques.

15 The term "alkyl C1 to 6" includes straight chain, branched, saturated,
unsaturated, aliphatic and cyclic alkyl groups containing 1 to 6 carbon atoms.

The compounds of formula I may exist in tautomeric, enantiomeric or
diastereomeric forms, all of which are included within the scope of the invention.

20 The various optical isomers may be isolated by separation of a racemic mixture of
the compounds using conventional techniques, e.g. fractional crystallisation, or
HPLC. Alternatively the individual enantiomers may be made by reaction of the
appropriate optically active starting materials under reaction conditions which will
not cause racemisation.

25

Intermediate compounds may also exist in enantiomeric forms and may be used
as purified enantiomers, diastereomers, racemates or mixtures.

The compounds of formula I possess useful pharmacological activity in animals.
30 In particular, they possess useful nitric oxide synthetase inhibiting activity, and are
expected to be useful in the treatment or prophylaxis of human diseases or

-12-

conditions in which the synthesis or oversynthesis of nitric oxide forms a contributory part; for example, hypoxia, e.g. in cases of cardiac arrest and stroke, neurodegenerative disorders including nerve degeneration and/or nerve necrosis in disorders such as hypoxia, hypoglycemia, epilepsy, and in external wounds (such as spinal cord and head injury), hyperbaric oxygen convulsions and toxicity, dementia e.g. pre-senile dementia, Alzheimer's disease and AIDS-related dementia, Sydenham's chorea, Parkinson's disease, Tourette's Syndrome, Huntington's disease, Amyotrophic Lateral Sclerosis, Korsakoff's disease. imbecility relating to a cerebral vessel disorder, sleeping disorders, schizophrenia, depression, autism, seasonal affective disorder, jet-lag, depression or other symptoms associated with Premenstrual Syndrome (PMS), anxiety and septic shock. Compounds of formula I may also be expected to show activity in the prevention and reversal of tolerance to opiates and diazepines, treatment of drug addiction, relief of pain and treatment of migraine and other vascular headaches. The compounds of the present invention may also show useful immunosuppressive activity, be useful in the treatment or prophylaxis of inflammation, neurogenic inflammation, reversible obstructive airways disease including asthma and adult respiratory distress syndrome (ARDS), in the treatment of gastrointestinal motility disorders, cancer, in the induction of labour, for reduction of gastric acid secretion and for increasing the contractile force of skeletal muscle.

Compounds of formula I are most particularly of interest in the treatment of neurodegenerative disorders, of migraine or for the prevention and reversal of tolerance to opiates and diazepines or for the treatment of drug addiction and especially in the treatment of neurodegenerative disorders.

Thus according to a further aspect of the invention we provide a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a pharmaceutical.

According to another feature of the invention we provide the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, in the

-13-

manufacture of a medicament for the treatment or prophylaxis of the
aforementioned diseases or conditions.

There is also provided a method of treatment or prophylaxis of one of the
5 aforementioned diseases or conditions which comprises administering a
therapeutically effective amount of a compound of formula I or a pharmaceutically
acceptable salt thereof, to a person suffering from or susceptible to such a disease
or condition.

10 For the above mentioned therapeutic indications, the dosage administered will,
of course, vary with the compound employed, the mode of administration and the
treatment desired. However, in general, satisfactory results are obtained when the
compounds are administered to a human at a daily dosage of between 1 mg and
2000 mg (measured as the solid form).

15 The compounds of formula I, and pharmaceutically acceptable salts thereof,
may be used on their own, or in the form of appropriate medicinal preparations for
enteral or parenteral administration.

20 According to the invention, there is provided a pharmaceutical formulation
including preferably less than 80% and more preferably less than 50% of a
compound of formula I, or a pharmaceutically acceptable salt thereof, in admixture
with a pharmaceutically acceptable diluent or carrier.

25 Examples of diluents and carriers which are suitable will be well known to a
person skilled in the art.

The enzyme nitric oxide synthetase has a number of isoforms and compounds of
formula I, or pharmaceutically acceptable salts thereof, may be screened for nitric
30 oxide synthetase activity by procedures based on those of Bredt and Snyder in Proc.
Natl. Acad. Sci. (1990) 87, 682-685 and Förstermann et. al., Eur. J. Pharm. (1992)

-14-

225, 161-165 as follows. Nitric oxide synthetase converts ^3H -L-arginine to ^3H -L-citrulline which can be separated by cation exchange chromatography and quantified by scintillation counting.

5 Screen A

(A) Screen for neuronal nitric oxide synthetase activity

Enzyme was isolated from rat hippocampus or cerebellum. The cerebellum or hippocampus of a male Sprague-Dawley rat (250-275g) is removed following CO_2 anaesthesia of the animal and decapitation. Cerebellar or hippocampal supernatant
10 is prepared by homogenisation in 50 mM Tris-HCl with 1 mM EDTA buffer (pH 7.2 at 25 °C) and centrifugation for 15 minutes at 20,000 g. Residual L-arginine is removed from the supernatant by chromatography through Dowex AG-50W-X8 sodium form and hydrogen form columns successively, and further centrifugation at 1000 g for 30 seconds.

15 For the assay, 25 μl of the final supernatant is added to each of 12 test tubes containing 25 μl L-arginine solution (of concentration 18 μM ^1H -L-arginine, 96 nM ^3H -L-arginine) and either 25 μl of an assay buffer (50 mM HEPES, 1 mM EDTA, 1.5 mM CaCl_2 , pH 7.4) or 25 μl of test compound in the buffer at 22 °C. To each test tube is added 75 μl of complete assay buffer (50 mM HEPES, 1 mM EDTA,
20 1.5 mM CaCl_2 , 1 mM DTT, 100 μM NADPH, 10 $\mu\text{g/ml}$ calmodulin, pH 7.4) to initiate the reaction and the reaction is stopped after 10 minutes by addition of 2 ml of a termination buffer (20 mM HEPES, 2 mM EDTA, pH 5.5).

Labelled L-citrulline is separated from labelled L-arginine by chromatography over a Dowex AG-50W-X8 200-400 mesh column. 1 ml of each terminated
25 reaction is added to an individual 1 ml column and the eluant combined with that from two 1 ml distilled water washes and 16 ml of scintillation cocktail. The L-citrulline is then quantified by scintillation counting.

In a typical experiment using the cerebellar supernatant, basal activity is increased by 20,000 dpm/ml of sample above a reagent blank which has an activity
30 of 7,000 dpm/ml. A reference standard, N-nitro-L-arginine, which gives 60%

-15-

inhibition of nitric oxide synthetase at a concentration of 1 μ M, is tested in the assay to verify the procedure.

Screen B

5 (B) Screen for macrophage nitric oxide synthetase activity

Enzyme is prepared, after induction, from the cultured murine macrophage cell line J774A-1 (obtained from laboratories of the Imperial Cancer Research Fund). J774A-1 cells are cultured in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% foetal bovine serum, 4 mM L-glutamine and antibiotics
10 (100 units/ml penicillin G, 100 μ g/ml streptomycin & 0.25 μ g/ml amphotericin B). Cells are routinely grown in 225 cm² flasks containing 35 ml medium kept at 37 °C and in a humidified atmosphere containing 5% CO₂.

Nitric oxide synthetase is produced by cells in response to interferon- γ (IFN γ) and lipopolysaccharide (LPS). The medium from confluent culture flasks is removed
15 and replaced with 25 ml (per flask) of fresh medium containing 1 μ g/ml LPS and 10 units/ml IFN γ . After a period of 17-20 hours in culture, harvesting of cells is accomplished by scraping the cell sheet from the flask surface into the culture medium. Cells are collected by centrifugation (1000g for 10 minutes) and lysate prepared by adding to the cell pellet a solution containing 50 mM Tris-HCl (pH 7.5
20 at 20 °C), 10% (v/v) glycerol, 0.1% (v/v) Triton-X-100, 0.1 μ M dithiothreitol and a cocktail of protease inhibitors comprising leupeptin (2 μ g/ml), soya bean trypsin inhibitor (10 μ g/ml), aprotinin (5 μ g/ml) & phenylmethylsulphonyl fluoride (50 μ g/ml).

For the assay, 25 μ l substrate cocktail (50 mM Tris-HCl (pH 7.5 at 20 °C), 400
25 μ M NADPH, 20 μ M flavin adenine dinucleotide, 20 μ M flavin mononucleotide, 4 μ M tetrahydrobiopterin, 12 μ M L-arginine and 0.025 μ Ci L-[³H] arginine) is added to wells of a 96 well filter plate (0.45 μ M pore size) containing 25 μ l of a solution of test compound in 50 mM Tris-HCl. The reaction is started by adding 50 μ l of cell lysate (prepared as above) and after incubation for 1 hour at room temperature is
30 terminated by addition of 50 μ l of an aqueous solution of 3 mM nitroarginine and 21 mM EDTA.

-16-

Labelled L-citrulline is separated from labelled L-arginine using Dowex AG-50W. 150 μ l of a 25% aqueous slurry of Dowex 50W (Na^+ form) is added to the assay after which the whole is filtered into 96 well plates. 70 μ l of filtrate is sampled and added to wells of 96 well plates containing solid scintillant. After
5 allowing the samples to dry the L-citrulline is quantified by scintillation counting.

In a typical experiment basal activity is 300 dpm per 70 μ l sample which is increased to 1900 dpm in the reagent controls. Aminoguanidine, which gives an IC_{50} (50% inhibitory concentration) of 10 μM , is tested as a standard to verify the procedure.

10

Screen C

(C) Screen for endothelial nitric oxide synthetase activity

Enzyme may be isolated from human umbilical vein endothelial cells (HUVECs) by a procedure based on that of Pollock *et al* (1991) Proc. Nat. Acad. Sci., 88,
15 10480-10484. HUVECs were purchased from Clonetics Corp (San Diego, CA, USA) and cultured to confluency. Cells can be maintained to passage 35-40 without significant loss of yield of nitric oxide synthetase. When cells reach confluency, they are resuspended in Dulbecco's phosphate buffered saline, centrifuged at 800 rpm for 10 mins, the cell pellet homogenised in ice-cold 50 mM
20 Tris-HCl, 1 mM EDTA, 10% glycerol, 1 mM phenylmethylsulphonylfluoride, 2 μM leupeptin at pH 4.2. Following centrifugation at 34,000 rpm for 60 mins. the pellet is solubilised in the homogenisation buffer which also contains 20 mM CHAPS. After a 30 min incubation on ice, the suspension is centrifuged at 34,000 rpm for 30 mins. The resulting supernatant is stored at -80 °C until use.

25 For the assay, 25 μ l of the final supernatant is added to each of 12 test tubes containing 25 μ l L-arginine solution (of concentration 12 μM ^1H -L-arginine, 64 nM ^3H -L-arginine) and either 25 μ l of an assay buffer (50 mM HEPES, 1 mM EDTA, 1.5 mM CaCl_2 , pH 7.4) or 25 μ l of test compound in the buffer at 22 °C. To each test tube was added 25 μ l of complete assay buffer (50 mM HEPES, 1 mM EDTA,
30 1.5 mM CaCl_2 , 1 mM DTT, 100 μM NADPH, 10 $\mu\text{g/ml}$ calmodulin, 12 μM tetrahydrobiopterin, pH 7.4) to initiate the reaction and the reaction is stopped

-17-

after 10 mins by addition of 2 ml of a termination buffer (20 mM HEPES, 2 mM EDTA, pH 5.5).

Labelled L-citrulline is separated from labelled L-arginine by chromatography over a Dowex AG-50W-X8 200-400 mesh column. 1 ml of each terminated
5 reaction is added to an individual 1 ml column and the eluant combined with that from two 1 ml distilled water washes and 16 ml of scintillation cocktail. The L-citrulline is then quantified by scintillation counting.

In a typical experiment, basal activity is increased by 5,000 dpm/ml of sample above a reagent blank which has an activity of 1500 dpm/ml. A reference standard,
10 N-nitro-L-arginine, which gives 70-90% inhibition of nitric oxide synthetase at a concentration of 1 μ M, is tested in the assay to verify the procedure.

Compounds may also be tested in an ex-vivo assay to determine the extent of brain penetration.

15 Screen D

(D) Ex vivo assay for neuronal nitric oxide synthetase activity

Male Sprague-Dawley rats (250-275g) were dosed intravenously at 10mg/kg with test compound dissolved in 0.9% saline or with saline alone as control. At a predetermined time (typically 2-24 hours) after treatment, the animals were
20 sacrificed, the cerebellum removed and the supernatant prepared and assayed for nitric oxide synthetase activity as described in Screen A.

As a further confirmatory test, a fraction of the cerebellar supernatant was applied to a 2'-5'-ADP Sepharose column (which binds nitric oxide synthetase) and subsequently eluted with NADPH. The eluant was tested for nitric oxide synthetase
25 activity following the procedure of Screen A.

Compounds that penetrate the rat brain and inhibit neuronal nitric oxide synthetase resulted in reduced nitric oxide synthetase activity both in the supernatant preparation and in the eluant from the 2'-5'-ADP Sepharose column.

30 In the screens for nitric oxide synthetase inhibition activity, compound activity is expressed as IC_{50} (the concentration of drug substance which gives 50% enzyme

-18-

inhibition in the assay). IC_{50} values for test compounds were initially estimated from the inhibiting activity of 1, 10 and 100 μ M solutions of the compounds. Compounds that inhibited the enzyme by at least 50% at 10 μ M were retested using more appropriate concentrations so that an IC_{50} could be determined.

5

In Screen A above (a screen for activity against the neuronal isoform of nitric oxide synthetase), the compound of Example 1 below gave an IC_{50} of less than 10 μ M indicating that it is expected to show useful therapeutic activity. In Screens B and C (the screens for activity against the macrophage and endothelial isoforms of nitric oxide synthetase) the compound of Example 1 gave IC_{50} values more than 10 times that obtained in Screen A indicating that it shows desirable selectivity.

The compounds of Examples 2-9, 10(a)-(f), 11-13 and 19-24 were also tested in Screen A and also gave IC_{50} values of less than 10 μ M. Thus these compounds are also expected to show useful therapeutic activity.

Compounds of formula I, and pharmaceutically acceptable salts thereof, have the advantage that they are less toxic, more efficacious, more selective, are longer acting, have a broader range of activity, are more potent, produce fewer side effects, are more easily absorbed, or have other useful pharmacological properties than compounds previously known and used in the therapeutic fields mentioned above.

Compounds of formula I, and pharmaceutically acceptable salts thereof, may also have the advantage that they are more selective for the neuronal isoform of nitric oxide synthetase enzyme and are therefore expected to show useful therapeutic activity with a reduced side-effect profile associated with inhibition of the other isoforms.

The invention is illustrated by the following examples:

Example 1

-19-

N-((2-(Phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate(a) 5-Nitro-2-indanone

This compound was prepared by the method of Heusler, Schieffer *Ber.*, (1899) 32, 33.

5 (b) 5-Nitro-2-(phenylmethyl)aminoindane

5-Nitro-2-indanone (1.48 g, 8.36 mmol), benzylamine (4.40 ml, 41.8 mmol), acetic acid (15.0 ml), 4 Å molecular sieves (20 ml), THF (15 ml), and MeOH (15 ml) were introduced into a flask and cooled to 0 °C. Sodium cyanoborohydride (1.05 g, 16.7 mmol) was then added portionwise over a 5-minute period. The mixture was stirred for 14 hr, filtered through celite, and concentrated to a syrup. The mixture was made basic with 2N NaOH and extracted with ether (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, concentrated and chromatographed over silica gel (3% methanol/methylene chloride) to yield 5-nitro-2-(phenylmethyl)aminoindane: (1.18 g, 53%); M.S. (M+H)⁺ = 269.

(c) 2-(5-Nitroindanyl)-N-(phenylmethyl)trifluoroacetamide

To a stirred solution of 5-nitro-2-(phenylmethyl)aminoindane (1.18 g, 4.40 mmol) and triethylamine (0.61 ml, 4.40 mmol) in methylene chloride (50 ml) was added trifluoroacetic anhydride (0.63 ml, 4.40 mmol) dropwise. After stirring for 1 minute, the solvent was dumped into water and extracted with methylene chloride (3 X 20 ml). The combined extracts were washed with water, dried over magnesium sulfate, and filtered through a short plug of silica gel (20% ethyl acetate/hexane) to yield 2-(5-nitroindanyl)-N-(phenylmethyl)trifluoroacetamide: (1.17g, 73%); M.S. (M+H)⁺ = 365.

25 (d) 2-(5-Aminoindanyl)-N-(phenylmethyl)trifluoroacetamide

To a stirred solution of 2-(5-nitroindanyl)-N-(phenylmethyl)trifluoroacetamide (1.17 g, 3.21 mmol) in THF/MeOH (100 ml, 1:1) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to give 2-(5-aminoindanyl)-N-(phenylmethyl)trifluoroacetamide which was homogeneous by TLC and used immediately in step (f).

(e) S-methyl-2-thiophenethiocarboximide hydroiodide

-20-

A solution of 2-thiophenecarboxthioamide (Maybridge Chemical) (11.1 g) in acetone (60 ml) was treated with iodomethane (13.4g). After 6 hrs at 22 °C, the resulting yellow solids were collected by filtration, washed with acetone (2 x 25ml) and dried to provide 18.45 g of S-methyl-2-thiophenethiocarboximide hydroiodide, m.p.195 °C (dec).

(f) N-((2-(Phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate
To a solution of 2-(5-aminoindanyl)-N-(phenylmethyl)trifluoroacetamide (1.0 g, 3.0 mmol) in isopropanol (6 ml)/DMF (0.5 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (0.85 g, 3.0 mmol). The mixture was stirred for 14 hr, diluted with methanol (6 ml) and 2 N NaOH (6 ml) and heated to 50 °C for 0.5 hr. The mixture was dumped into water and extracted with ethyl acetate (3 X 30 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and chromatographed over silica gel (20% methanol/methylene chloride) to give the titled compound as the free base. Treatment with IPA/oxalic acid yielded N-((2-(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate as a white solid: (0.47 g, 30%); m.p. 130-135 °C.

Example 2

N-((2-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) 7-Nitro-3,4-dihydro-2(1H)-naphthaleneone

This compound was prepared following the method of J. Med. Chem. (1989) 32, 2128.

(b) 7-Nitro-2-((phenylmethyl)amino)-1,2,3,4-tetrahydronaphthalene

7-Nitro-3,4-dihydro-2(1H)-naphthaleneone (1.50 g, 7.85 mmol), benzylamine (4.30 ml, 39.3 mmol), acetic acid (8.0 ml), 4 Å molecular sieves (20 ml), THF (15 ml), and MeOH (15 ml) were introduced into a flask and cooled to 0 °C. Sodium cyanoborohydride (0.99 g, 15.7 mmol) was then added portionwise over a 5-minute period. The mixture was stirred for 14 hr, filtered through celite, and concentrated to a syrup. The mixture was made basic with 2N NaOH and extracted with ether (3 X 50 ml). The combined extracts were washed with water, dried over magnesium

-21-

sulfate, filtered, concentrated and chromatographed over silica gel (3% methanol/methylene chloride) to yield 7-nitro-2-((phenylmethyl)amino)-1,2,3,4-tetrahydronaphthalene: (2.10 g, 95%); M.S. (M+H)⁺ = 283.

(c) 2-(7-Nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide

- 5 To a stirred solution of 7-nitro-2-((phenylmethyl)amino)-1,2,3,4-tetrahydronaphthalene (2.10 g, 7.45 mmol) and triethylamine (1.07 ml, 7.45 mmol) in methylene chloride (50 ml) was added trifluoroacetic anhydride (1.05 ml, 7.45 mmol) dropwise. After stirring for 1 minute, the solvent was dumped into water and extracted with methylene chloride (3 X 20 ml). The combined extracts were
10 washed with water, dried over magnesium sulfate, and filtered through a short plug of silica gel (20% ethyl acetate/hexane) to yield 2-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide: (2.55 g, 90%); M.S. (M+H)⁺ = 379.

(d) 2-(7-Amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide

- 15 To a stirred solution of 2-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide (2.55 g, 6.75 mmol) in THF/MeOH (100 ml, 1:1) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to give 2-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide which was homogeneous
20 by TLC and used immediately in the next step.

(e) N-((2-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

- To a solution of 2-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide (2.11 g, 6.07 mmol) in isopropanol (10 ml) was
25 added S-methyl-2-thiophenethiocarboximide hydroiodide (1.72 g, 6.07 mmol). The mixture was stirred for 14 hr, diluted with methanol (6 ml) and 2 N NaOH (6 ml) and heated to 50 °C for 0.5 hr. The mixture was dumped into water and extracted with ethyl acetate (3 X 30 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and concentrated to a solid which was
30 recrystallized (methylene chloride/hexane) to yield N-((2-(phenylmethyl)amino)-

-22-

1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide as a white solid: (0.66 g, 30%); m.p. 119-120 °C.

Example 3

5 N-((2-Amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dioxalate

(a) 7-Nitro-2-amino-1,2,3,4-tetrahydronaphthalene hydrochloride

7-Nitro-1-tetralone (1.50 g, 7.85 mmol), ammonium acetate (6.05 ml, 78.5 mmol), acetic acid (8.0 ml), 4 Å molecular sieves (20 ml), THF (15 ml) and MeOH (15 ml) were introduced into a flask and cooled to 0 °C. Sodium cyanoborohydride (0.99 g, 10 15.7 mmol) was then added portionwise over a 5-minute period. The mixture was stirred for 1 hr, filtered through celite, and concentrated to a syrup. The mixture was made basic with 2N NaOH and extracted with ether (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and concentrated to yield an oil. The compound was isolated as the hydrochloride 15 salt : 7-nitro-2-amino-1,2,3,4-tetrahydronaphthalene hydrochloride: (1.00 g, 56%); m.p. >300 °C.

(b) 2-(7-Nitro-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide

To a stirred solution of 7-nitro-2-amino-1,2,3,4-tetrahydronaphthalene hydrochloride (1.00 g, 4.39 mmol) and triethylamine (1.22 ml, 8.77 mmol) in 20 methylene chloride (50 ml) was added trifluoroacetic anhydride (0.62 ml, 4.39 mmol) dropwise. After stirring for 1 minute, the solvent was dumped into water and extracted with methylene chloride (3 X 20 ml). The combined extracts were washed with water, dried over magnesium sulfate, and filtered through a short plug of silica gel (20% ethyl acetate/hexane) to yield 2-(7-nitro-(1,2,3,4- 25 tetrahydronaphthyl))-N-trifluoroacetamide: (0.78 g, 62%); M.S. (M+H)⁺ = 289.

(c) 2-(7-Amino-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide

To a stirred solution of 2-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide (0.76 g, 2.21 mmol) in THF/MeOH (100 ml, 1:1) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, 30 and concentrated to give 2-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-

-23-

trifluoroacetamide which was homogeneous by TLC and used immediately in the next reaction.

(d) N-((2-Amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dihydrobromide

- 5 To a solution of 2-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide (0.70 g, 2.71 mmol) in isopropanol (10 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (0.77 g, 2.71 mmol). The mixture was stirred for 14 hr, diluted with methanol (6 ml) and 2 N NaOH (6 ml) and heated to 50 °C for 0.5 hr. The mixture was dumped into water and extracted with ethyl acetate (3
10 X 30 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, concentrated, and chromatographed over silica (20 % methanol/methylene chloride) to yield an oil which was converted to the dihydrobromide salt: N-((2-amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dihydrobromide: (0.37 g, 32%); dec >210 °C.

15

Example 4

N-((1-Amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dioxalate

(a) 7-Nitro-1-amino-1,2,3,4-tetrahydronaphthalene

- 7-Nitro-1-amino-1,2,3,4-tetrahydronaphthalene was made as for 7-Nitro-2-amino-
20 1,2,3,4-tetrahydronaphthalene. The compound was isolated as the hydrochloride salt : (0.30 g, 12%); m.p. >300 °C.

(b) 1-(7-Nitro-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide

1-(7-Nitro-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide was made as for 2-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide: (0.35 g, 95%); M.S.

- 25 (M+H)⁺ = 289.

(c) 1-(7-Amino-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide

1-(7-Amino-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide was made as for 2-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-trifluoroacetamide and used immediately in the next reaction.

- 30 (d) N-((1-Amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dioxalate

-24-

N-((1-Amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dioxalate was made as for N-((2-amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dihydrobromide except that it was isolated as the dioxalate salt: (0.18 g, 33%); dec >155 °C.

5

Example 5N-((2-Amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate(a) 5-Nitro-2-aminoindane hydrochloride

To 2-aminoindane hydrochloride (19.11 g, 0.112 mol) at 0 °C was added sulfuric acid (60 ml) followed by potassium nitrate (11.84 g, 0.117 mol). The mixture was allowed to warm to room temperature, stirred for an additional 2 hr, then dumped onto ice/50% NaOH (500 ml total). The mixture was extracted with ether (3 X 200 ml) and the combined extracts were washed with water, dried over magnesium sulfate, filtered, and concentrated to an oil which was converted to the hydrochloride salt. Recrystallization from isopropanol/methanol afforded 5-nitro-2-aminoindane hydrochloride: (14.58 g, 60%); m.p. >300 °C.

(b) 2-(5-Nitroindanyl)-N-trifluoroacetamide

To a stirred solution of 5-nitro-2-aminoindane hydrochloride (1.00 g, 5.89 mmol) and triethylamine (0.82ml, 5.89 mmol) in methylene chloride (50 ml) was added trifluoroacetic anhydride (0.83 ml, 5.89 mmol) dropwise. After stirring for 1 minute, the solvent was dumped into water and extracted with methylene chloride (3 X 20 ml). The combined extracts were washed with water, dried over magnesium sulfate, and filtered through a short plug of silica gel (20% ethyl acetate/hexane) to yield 2-(5-nitroindanyl)-N-trifluoroacetamide: (1.51 g, 93%); m.p. 153-154 °C.

(c) 2-(5-Aminoindanyl)-N-trifluoroacetamide

To a stirred solution of 2-(5-nitroindanyl)-N-trifluoroacetamide (0.58 g, 2.25 mmol) in THF/MeOH (100 ml, 1:1) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to give 2-(5-aminoindanyl)-N-trifluoroacetamide which was homogeneous by TLC and used immediately in the next step.

(d) N-((2-Amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate

30

-25-

To a solution of 2-(5-aminoindanyl)-N-trifluoroacetamide (0.52 g, 2.25 mmol) in isopropanol (6 ml)/DMF (0.5 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (0.64 g, 2.25 mmol). The mixture was stirred for 14 hr, diluted with methanol (6 ml) and 2 N NaOH (6 ml) and heated to 50 °C for 0.5 hr. The
5 mixture was dumped into water and extracted with ethyl acetate (3 X 30 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and chromatographed over silica gel (20% methanol/methylene chloride) to give the titled compound as the free base. Treatment with IPA/oxalic acid yielded N-((2-amino)indan-5-yl)-2-thiophenecarboximidamide dioxalate as a white solid: (0.60
10 g, 50%); m.p. dec 70 °C.

Example 6

N-((2-(Methyl)(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dihydrobromide

15 (a) 5-Nitro-2-(phenylmethyl)aminoindane hydrochloride

To 5-nitro-2-aminoindane hydrochloride (3.00 g, 14.00 mmol) in DMF (60 ml) was added triethylamine (4.07 ml, 29.40 mmol) followed by benzyl bromide (1.74 ml, 14.68 mmol). The mixture was warmed to room temperature, stirred for 1 hr, dumped into water (200 ml), and extracted with ethyl acetate (3 X 50 ml). The
20 combined extracts were washed with water, dried over magnesium sulfate, filtered through a small plug of silica gel and reduced to a syrup. The sub-titled compound was isolated as the hydrochloride salt: (2.29 g, 54%); m.p. dec 266 °C.

(b) 5-Nitro-2-(methyl)(phenylmethyl)aminoindane hydrochloride

To 5-nitro-2-(phenylmethyl)aminoindane hydrochloride (2.29 g, 7.52 mmol) in DMF
25 (100 ml) was added potassium carbonate (2.60 g, 18.80 mmol) followed by methyl iodide (0.47 ml, 7.52 mmol). The mixture was warmed to room temperature, stirred for 16 hr, dumped into water (400 ml), and extracted with ethyl acetate (3 X 100 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered through a small plug of silica gel and reduced to a syrup. The titled
30 compound was isolated as the hydrochloride salt: (1.08 g, 45%); m.p. dec 280 °C.

(c) 5-Amino-2-(methyl)(phenylmethyl)aminoindane dihydrochloride

-26-

To 5-nitro-2-(methyl)(phenylmethyl)aminoindane hydrochloride (1.08 g, 3.39 mmol) in 85% acetic acid/water was added zinc powder (3.0 g). The mixture was stirred for 1 minute, filtered through celite and concentrated. The concentrate was neutralized with 2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, and reduced to a syrup. The oil was treated with IPA/HCl, concentrated and used immediately in the next step.

(d) N-((2-(Methyl)(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dihydrobromide

To 5-amino-2-(methyl)(phenylmethyl)aminoindane dihydrochloride in DMF (10 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (0.98 g, 3.45 mmol) and pyridine (0.27 ml, 3.29 mmol). The mixture was stirred for 14 hr, dumped onto water/2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and chromatographed over silica gel (10% methanol/methylene chloride) to give the titled compound as the free base. Treatment with IPA/HBr yielded N-((2-(methyl)(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide dihydrobromide as a white solid: (0.43 g, 25%); m.p. 196-200 °C.

Example 7

N-((1-amino)indan-6-yl)-2-thiophenecarboximidamide dihydrochloride

(a) 6-Nitro-1-aminoindane hydrochloride

1-Aminoindane (10.0 g, 75.08 mmol) was added to concentrated sulfuric acid (40 ml) at 0 °C. The mix was warmed to room temperature to aid in solvation then cooled to 0 °C. Potassium nitrate (7.60 g, 75.08 mmol) was then added portionwise and the mixture allowed to stir at room temperature for 1 hr before being dumped onto ice/50 % NaOH. The aqueous solution was extracted with chloroform (3 X 100 ml). The combined extracts were washed with water, decolorized with charcoal, dried over magnesium sulfate, filtered, and concentrated to an oil. Treatment with IPA/HCl afforded the sub-titled compound: (6.90 g, 43%); m.p. dec 280 °C.

(b) 6-Amino-1-aminoindane hydrochloride

-27-

To a solution of 6-nitro-1-aminoindane hydrochloride (1.00 g, 4.66 mmol) in MeOH (100 ml) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to give 6-amino-1-aminoindane hydrochloride which was homogeneous by TLC and used immediately in the next step.

(c) N-((1-amino)indan-6-yl)-2-thiophenecarboximidamide dihydrochloride

To 6-amino-1-aminoindane hydrochloride (0.74 g, 4.01 mmol) in DMF/IPA (4 ml, 1:1) was added S-methyl-2-thiophenethiocarboximide hydroiodide (1.26 g, 4.41 mmol). The mixture was heated to 50 °C, stirred for 16 hr then dumped into water/2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to an oil. Treatment with IPA/HCl yielded N-((1-amino)indan-6-yl)-2-thiophenecarboximidamide dihydrochloride as a white solid: (0.79 g, 60%); m.p. dec > 200 °C.

Example 8

N-((1-(Phenylmethyl)amino)indan-6-yl)-2-thiophenecarboximidamide dioxalate

(a) 6-Nitro-1-(phenylmethyl)aminoindane hydrochloride

To 6-nitro-1-aminoindane hydrochloride (1.90 g, 8.85 mmol) in DMF (30 ml) was added triethylamine (2.50 ml, 18.06 mmol) followed by benzyl bromide (1.07 ml, 9.03 mmol). The mixture was warmed to room temperature, stirred for 3 hr, dumped into water (100 ml), and extracted with ethyl acetate (3 X 70 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered through a small plug of silica gel and reduced to a syrup. The titled compound was isolated as the hydrochloride salt: (1.34 g, 50%); m.p. 234-235 °C.

(b) 6-Amino-1-(phenylmethyl)aminoindane hydrochloride

To 6-nitro-1-(phenylmethyl)aminoindane hydrochloride (1.34 g, 4.40 mmol) in MeOH (100 ml) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to give 6-amino-1-(phenylmethyl)aminoindane hydrochloride which was homogeneous by TLC and used immediately in the next step.

-28-

(c) N-((1-(Phenylmethyl)amino)indan-6-yl)-2-thiophenecarboximidamide dioxalate

To 6-amino-1-(phenylmethyl)aminoindane hydrochloride (1.21 g, 4.40 mmol) in DMF (20 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (1.38 g, 4.84 mmol). The mixture was heated to 50 °C, stirred for 16 hr then dumped into
5 water/2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to an oil. Treatment with IPA/oxalic acid yielded N-((1-(phenylmethyl)amino)indan-6-yl)-2-thiophenecarboximidamide dioxalate as a white solid: (1.06 g, 46%); m.p. dec > 120 °C.

10

Example 9

N-((2-((3-Chlorophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboximidine

(a) 2-((3-Chlorophenyl)carbonyl)amino-6-nitroindane

To 2-amino-6-nitroindane hydrochloride (1.5 g, 7.0 mmol) in methylene chloride (50
15 ml) at 0 °C was added triethylamine (2.1 ml, 15.0 mmol) followed by 3-chlorobenzoyl chloride (1.0 ml, 7.5 mmol). The mixture was dumped immediately into water and the layers separated. The aqueous layer was extracted with methylene chloride (2 X 20 ml) and the combined extracts washed with water, dried over MgSO₄, filtered, and concentrated to an oil which was homogeneous by TLC and used immediately
20 in the next step: M.S. (M+H)⁺ = 317.

(b) 2-((3-Chlorophenyl)methyl)amino-6-nitroindane

To 2-((3-chlorophenyl)carbonyl)amino-6-nitroindane (2.2 g, 7.0 mmol) in THF (75 ml) was added BH₃•THF (1.0 M, 35 ml, 35 mmol) dropwise. The mixture was refluxed for 12 hr, cooled to 0 °C, quenched with 4N HCl (60 ml), and refluxed for
25 1 hr. The resulting solution was evaporated to an oil, made basic with 50% NaOH, and extracted with methylene chloride (3 X 20 ml). The combined extracts were washed with water, dried over MgSO₄, filtered and concentrated to an oil.

Treatment with IPA/HCl yielded 2-((3-chlorophenyl)methyl)amino-6-nitroindane: (2.1 g, 88% two steps); m.p. 234-237 °C.

30 (c) 2-((3-chlorophenyl)methyl)amino-6-aminoindane

-29-

To 2-((3-chlorophenyl)methyl)amino-6-nitroindane hydrochloride (2.1 g, 6.13 mmol) in 85% AcOH/H₂O (40 ml) was added zinc metal (1.6 g, 24.5 mmol). The mixture was stirred for 5 min, filtered through celite, and evaporated to an oil. The oil was dumped into basic water and extracted with chloroform (3 X 20 ml). The combined extracts were washed with water, dried over MgSO₄, filtered and concentrated to an oil. Treatment with IPA/HCl yielded 2-((3-chlorophenyl)methyl)amino-6-aminoindane: (1.5 g, 70%); m.p. > 270 °C.

- (d) N-((2-((3-Chlorophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine
2-((3-Chlorophenyl)methyl)amino-6-aminoindane dihydrochloride (1.5 g, 4.2 mmol),
10 S-methyl-2-thiophenethiocarboximide hydroiodide (1.3 g, 4.6 mmol) and pyridine (0.34 ml, 4.2 mmol) in DMF (10 ml) were stirred for 24 hr. The mixture was dumped into water, made basic with 2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over MgSO₄, filtered, concentrated, and chromatographed over silica gel (12% MeOH/methylene chloride) to afford a colorless oil. Treatment with IPA/HCl yielded N-((2-((3-chlorophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine: (0.75 g, 40%);
15 m.p. 297-299 °C.

Example 10

- 20 The following compounds were prepared according to the method of Example 9:
- (a) N-((2-((2-Methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 183 °C.
- (b) N-((2-((3-Methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 195 °C.
- 25 (c) N-((2-((4-Methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 182 °C.
- (d) N-((2-(Ethylamino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 236-238 °C.
- (e) N-((2-(((4-Phenyl)phenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 182 °C.
- 30 (f) N-((2-(((4-Hexyl)phenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine; m.p. 125 °C.

-30-

(g) N-((2-((3-Bromophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboximidine; m.p. 182 °C.

Example 11

5 N-((2-((3-Chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) 7-Nitro-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene

7-Nitro-3,4-dihydro-2(1H)-naphthaleneone (1.50 g, 7.85 mmol), 3-chlorobenzylamine (4.70 ml, 39.3 mmol), acetic acid (6.0 ml), 4 Å molecular sieves (20 ml), THF (15 ml), and MeOH (15 ml) were introduced into a flask and cooled to 0 °C. Sodium cyanoborohydride (0.99 g, 15.7 mmol) was added portionwise over a 5-minute period. The mixture was stirred for 14 hr, filtered through celite, and concentrated to a syrup. The mixture was made basic with 2N NaOH and extracted with ether (3 X 50 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, concentrated and chromatographed over silica gel (3% methanol/methylene chloride). Treatment of the oil with IPA/HCl yielded 7-nitro-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene hydrochloride: (1.34 g, 50%); M.S. (M+H)⁺ = 317.

(b) 7-Amino-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene

20 To 7-nitro-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene hydrochloride (1.34 g, 3.80 mmol) in 85% AcOH/H₂O (75 ml) was added zinc metal (2.48 g, 38.0 mmol). The mixture was stirred for 5 min, filtered through celite, and evaporated to an oil. The oil was dumped into basic water and extracted with chloroform (3 X 20 ml). The combined extracts were washed with water, dried over MgSO₄, filtered and concentrated to an oil. Treatment with IPA/HCl yielded 7-amino-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene (1.4 g, 99%); M.S. (M+H)⁺ = 288.

(c) N-((2-((3-Chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

30 7-Amino-2-(((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphthalene dihydrochloride (1.32 g, 3.70 mmol), S-methyl-2-thiophenethiocarboximide

-31-

hydroiodide (1.3 g, 4.6 mmol), and pyridine (0.30 ml, 3.7 mmol) in DMF (15 ml) were stirred for 24 hr. The mixture was dumped into water, made basic with 2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over MgSO_4 , filtered and concentrated to an oil.

- 5 Treatment with IPA/oxalic acid yielded N-((2-((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dioxalate: (0.71 g, 33%); dec > 100 °C.

Example 12

- 10 N-((2-(Phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) 7-Nitro-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene

- To a stirred solution of 7-nitro-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene (1.5 g, 5.4 mmol) in DMF (30 ml) was added potassium
15 carbonate (1.5 g, 10.8 mmol) and methyl iodide (0.36 ml, 5.8 mmol). The mixture was stirred for 24 hr, dumped into water and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over MgSO_4 , filtered and concentrated to an oil. Treatment with IPA/HCl yielded 7-nitro-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene hydrochloride: (0.89
20 g, 50%); M.S. $(\text{M}+\text{H})^+ = 297$.

(b) 7-Amino-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene

- To 7-nitro-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene hydrochloride (0.89 g, 2.7 mmol) in 85% $\text{AcOH}/\text{H}_2\text{O}$ (75 ml) was added zinc metal (3.5 g, 54.0 mmol). The mixture was stirred for 5 min, filtered through celite, and
25 evaporated to an oil. The oil was dumped into basic water and extracted with chloroform (3 X 20 ml). The combined extracts were washed with water, dried over MgSO_4 , filtered and concentrated to an oil. Treatment with IPA/HCl yielded 7-amino-2-((phenyl)methyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene: (0.81 g, 88%); M.S. $(\text{M}+\text{H})^+ = 267$.

- 30 (c) N-((2-(Phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

-32-

7-Amino-2-((phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphthalene dihydrochloride (0.81 g, 2.4 mmol), S-methyl-2-thiophenethiocarboximide hydroiodide (0.74 g, 2.6 mmol) and pyridine (0.19 ml, 2.4 mmol) in DMF (15 ml) were stirred for 24 hr. The mixture was dumped into water, made basic with 2N NaOH and extracted with ethyl acetate (3 X 50 ml). The combined extracts were washed with water, dried over MgSO_4 , filtered, concentrated and chromatographed over silica gel (15% MeOH/methylene chloride). Concentration of the fractions yielded a solid which was recrystallized from ethyl acetate/hexane affording N-((2-(phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dihydrochloride: (0.14 g, 16%); m.p 176-178 °C.

Example 13

N-((1-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) 7-Nitro-1-((phenylmethyl)amino)-1,2,3,4-tetrahydronaphthalene
7-Nitro-1-tetralone (2.0 g, 10.5 mmol), benzylamine (1.2 ml, 10.5 mmol) and titanium isopropoxide (3.9 ml, 13.1 mmol) were combined and stirred for 1 hr. The mixture was diluted with absolute ethanol (12 ml), treated with sodium cyanoborohydride (0.44 g, 7.0 mmol) and allowed to stir for 20 hr. The solids were filtered and washed with ethanol. The ethanol was concentrated and the remaining oil used immediately in the next reaction: M.S. $(\text{M}+\text{H})^+ = 283$.

(b) 1-(7-Nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide

To a stirred solution of 7-nitro-1-((phenylmethyl)amino)-1,2,3,4-tetrahydronaphthalene (2.96 g, 10.50 mmol) and triethylamine (1.46 ml, 10.50 mmol) in methylene chloride (50 ml) was added trifluoroacetic anhydride (1.46 ml, 10.50 mmol) dropwise. After stirring for 1 minute, the solvent was dumped into water and extracted with methylene chloride (3 X 20 ml). The combined extracts were washed with water, dried over magnesium sulfate, and filtered through a short plug of silica gel (20% ethyl acetate/hexane) to yield 1-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide: (1.90 g, 48% two steps); M.S. $(\text{M}+\text{H})^+ = 379$.

-33-

(c) 1-(7-Amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide

To a stirred solution of 1-(7-nitro-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide (1.91 g, 5.05 mmol) in THF/MeOH (100 ml, 1:1) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50
5 psi for 1 hr, filtered through celite, and concentrated to give 1-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide which was homogeneous by TLC and used immediately in the next step.

(d) N-((1-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

10 To a solution of 1-(7-amino-(1,2,3,4-tetrahydronaphthyl))-N-(phenylmethyl)trifluoroacetamide (1.76 g, 5.05 mmol) in isopropanol (10 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (1.44 g, 5.05 mmol). The mixture was stirred for 14 hr, diluted with methanol (6 ml) and 2 N NaOH (6 ml) and heated to 50 °C for 0.5 hr. The mixture was dumped into water and extracted
15 with ethyl acetate (3 X 30 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and concentrated to an oil. Treatment with IPA/HBr yielded N-((2-(phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide dihydrobromide as a white solid: (0.53 g, 20%); m.p. 260-262 °C.

20

Example 14

N-((1-(Phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide

(a) 6-Acetamido-1-((phenyl)methyl)amino)indane

6-Acetamido-1-indanone (5.0 g, 27.6 mmol), benzylamine (3.1 ml, 27.9 mmol), and
25 titanium isopropoxide (10.2 ml, 34.5 mmol) were combined and stirred for 1 hr. The mixture was diluted with absolute ethanol (30 ml), treated with sodium cyanoborohydride (1.2 g, 19.3 mmol) and allowed to stir for 20 hr. The solids were filtered and washed with ethanol. The ethanol was concentrated and the remaining oil dissolved in ethyl acetate and extracted with 1N HCl (3 X 50 ml). The aqueous
30 layer was neutralized with 2N NaOH and extracted with ethyl acetate (3 X 100 ml). The combined extracts were washed with water, dried over magnesium sulfate,

-34-

filtered, and concentrated to an oil which was used without purification in the next step.

(b) 6-Amino-(1-((phenyl)methyl)amino)indane

6-Acetamido-(1-((phenyl)methyl)amino)indane was refluxed in 4N HCl (50 ml) for
5 20 min, cooled and extracted with ethyl acetate (3 X 50 ml). The aqueous layer
was neutralized with 2N NaOH and extracted with ethyl acetate (3 X 100 ml).
The combined extracts were washed with water, dried over magnesium sulfate,
filtered and concentrated to an oil. The oil was dissolved in IPA and treated with
IPA/HCl yielding the dihydrochloride salt: (2.0g, 24% for two steps); m.p. dec >
10 250 °C.

(c) N-((1-(Phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide

To 6-amino-(1-((phenyl)methyl)amino)indane dihydrochloride (2.0 g, 6.4 mmol) in
DMF (20 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (2.2 g,
7.7 mmol) and pyridine (0.57 ml, 7.1 mmol). The mixture was stirred at 50 °C for 20
15 hr, dumped into basic water and extracted with ethyl acetate (3 X 100 ml). The
combined extracts were washed with water, dried over magnesium sulfate, filtered,
concentrated and chromatographed over silica gel (6% methanol/methylene
chloride). The extracts were concentrated to an oil which was dissolved in
methanol, treated with IPA/HCl and triturated with ether. The solids were
20 collected by filtration and washed with ether: (1.1 g. 40%); m.p. dec > 180 °C.

Example 15

The following compound was prepared following the method of Example 14:

N-((1-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-6-yl)-2-
25 thiophenecarboximidamide m.p. dec > 200 °C.

Example 16

N-((2-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-furancarboximidamide

(a) 2-((Phenyl)carbonyl)amino-7-nitrotetralin

30 To 2-amino-7-nitrotetralin (2.8 g, 14.5 mmol) in THF (50 ml) and 10% K₂CO₃ (100
ml) was added benzoyl chloride (1.7 ml, 15.3 mmol). After the addition was

-35-

complete, the mixture was diluted with water to a volume of 250 ml. The precipitated solids were collected by filtration, washed with water, and dried *in vacuo*: (4.2 g, 98%), m.p. 194-198 °C.

(b) 2-((Phenyl)methyl)amino-7-nitrotetralin hydrochloride

- 5 To 2-((phenyl)carbonyl)amino-7-nitrotetralin (4.2 g, 14.1 mmol) in anhydrous THF (100 ml) was added borane-THF (49.3 ml, 1M THF, 49.3 mmol). The mixture was refluxed for 5 hr, cooled to 0 °C, and quenched with the dropwise addition of 4N HCl. The mixture was again brought to reflux for 1 hr, concentrated in *vacuo*, and the solids filtered (washed with water) and dried *in vacuo*: (3.5 g, 78%), m.p. > 300
10 °C.

(c) 2-((Phenyl)methyl)amino-7-aminotetralin hydrochloride

- To a stirred solution of 2-((phenyl)methyl)amino-7-nitrotetralin (2.0 g, 6.3 mmol) in MeOH (100 ml) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to an oil
15 which was homogeneous by TLC and used immediately in the next step.

(d) N-((2-((Phenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-furancarboximidamide

- To 2-((phenyl)methyl)amino-7-aminotetralin hydrochloride (1.8 g, 6.3 mmol) in DMF (20 ml) was added S-methyl-2-furanthiocarboximide hydroiodide (2.0 g, 7.5
20 mmol). The mixture was stirred for 2 hr at 45 °C, dumped into basic water and extracted with ethyl acetate (3 X 100 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to an oil. The oil was dissolved in methanol, treated with IPA/HCl and triturated with ether. The solids were collected by filtration and washed with ether: (2.2 g, 84%), m.p. dec >
25 195 °C.

Preparation of chiral intermediate compounds for Examples 17 and 18

Resolution of 2-amino-7-nitrotetralin

- 2-Amino-7-nitrotetralin (30 g, 156 mmol) dissolved in 200 ml of acetone was added
30 to dibenzoyl-D-tartaric acid (58.7 g, 164 mmol) also dissolved in 200 ml of acetone. The thick paste was filtered and washed with acetone. The paste was refluxed in

-36-

3L of water/ethanol/acetonitrile (1:1:1) then filtered hot. The solids collected upon filtration were recrystallized from the above mixture (3X): (5.25 g, 6%) of a single isomer was obtained as determined by chiral capillary zone electrophoresis, m.p. 240-242 °C.

5

Likewise, dibenzoyl-L-tartaric acid could be employed to resolve the opposite enantiomer using the same solvent system as that described above: (5.3 g, 6%), of a single isomer was obtained as determined by chiral capillary zone electrophoresis, m.p. 240-242 °C.

10

Example 17

(+)-N-((2-((Phenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) (+)-2-((Phenyl)carbonyl)amino-7-nitrotetralin

15 To 2-amino-7-nitrotetralin (1.8 g, 9.39 mmol, derived from dibenzoyl-D-tartaric acid) in THF (50 ml) and 10% K₂CO₃ (100 ml) was added benzoyl chloride (1.2 ml, 10.1 mmol). After the addition was complete, the mixture was diluted with water to a volume of 250 ml. The precipitated solids were collected by filtration, washed with water and dried *in vacuo*: (2.8 g, 100%), m.p. 208-209 °C, [α]_D +21.9 ° (c 0.33 DMSO).

20

(b) (+)-2-((phenyl)methyl)amino-7-nitrotetralin hydrochloride

To (+)-2-((phenyl)carbonyl)amino-7-nitrotetralin (2.8 g, 9.4 mmol) in anhydrous THF (100 ml) was added borane-THF (32.8 ml, 1M THF, 32.8 mmol). The mixture was refluxed for 5 hr, cooled to 0 °C, and quenched with the dropwise addition of 25 4N HCl. The mixture was brought to reflux for 1 hr, concentrated in vacuo, and the solids filtered (washed with water) and dried *in vacuo*: (2.8 g, 94%), m.p. > 300 °C, [α]_D +51.0 ° (c 0.33 DMSO).

(c) (+)-2-((Phenyl)methyl)amino-7-aminotetralin hydrochloride

To a stirred solution of (+)-2-((phenyl)methyl)amino-7-nitrotetralin (2.8 g, 8.7 mmol) in MeOH (100 ml) was added a catalytic amount of 10% Pd/C. The mixture 30

-37-

was hydrogenated at 50 psi for 1 hr, filtered through celite, and concentrated to a glassy solid which was homogeneous by TLC; $[\alpha]_D +73.3^\circ$ (c 0.87 DMSO).

(d) (+)-N-((2-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

- 5 To (+)-2-((phenyl)methyl)amino-7-aminotetralin hydrochloride (2.5 g, 8.7 mmol) in DMF (20 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (3.0 g, 10.4 mmol). The mixture was stirred for 4 hr at 45 °C, dumped into basic water and extracted with ethyl acetate (3 X 100 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered, and concentrated to an oil. The oil was dissolved in methanol, treated with IPA/HCl and triturated with ether. The solids were collected by filtration and washed with ether. One recrystallization from IPA/MeOH/Et₂O yielded a white solid: (2.5 g, 66%), m.p. dec > 260 °C, $[\alpha]_D +44.5^\circ$ (c 0.62 DMSO).
- 10

15 Example 18

(-)-N-((2-((Phenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

(a) (-)-2-((Phenyl)carbonyl)amino-7-nitrotetralin

- To 2-amino-7-nitrotetralin (1.8 g, 9.39 mmol, derived from dibenzoyl-L-tartaric acid) in THF (50 ml) and 10% K₂CO₃ (100 ml) was added benzoyl chloride (1.2 ml, 10.1 mmol). After the addition was complete, the mixture was diluted with water to a volume of 250 ml. The precipitated solids were collected by filtration, washed with water and dried *in vacuo*: (2.8 g, 100%), m.p. 208-209 °C, $[\alpha]_D -24.0^\circ$ (c 0.87 DMSO).
- 20

25 (b) (-)-2-((phenyl)methyl)amino-7-nitrotetralin hydrochloride

- To (-)-2-((phenyl)carbonyl)amino-7-nitrotetralin (2.8 g, 9.4 mmol) in anhydrous THF (100 ml) was added borane-THF (32.8 ml, 1M THF, 32.8 mmol). The mixture was refluxed for 5 hr, cooled to 0 °C, and quenched with the dropwise addition of 4N HCl. The mixture was again brought to reflux for 1 hr, concentrated *in vacuo*, and the solids filtered (washed with water) and dried *in vacuo*: (2.8 g, 94%), m.p. > 300 °C, $[\alpha]_D -59.4^\circ$ (c 0.39 DMSO).
- 30

-38-

(c) (-)-2-((Phenyl)methyl)amino-7-aminotetralin hydrochloride

To a stirred solution of (-)-2-((phenyl)methyl)amino-7-nitrotetralin (2.8 g, 8.7 mmol) in MeOH (100 ml) was added a catalytic amount of 10% Pd/C. The mixture was hydrogenated at 50 psi for 1 hr, filtered through celite and concentrated to a glassy solid which was homogeneous by TLC; $[\alpha]_D -74.6^\circ$ (c 0.80 DMSO).

(d) (-)-N-((2-(Phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide

To (-)-2-((phenyl)methyl)amino-7-aminotetralin hydrochloride (2.5 g, 8.7 mmol) in DMF (20 ml) was added S-methyl-2-thiophenethiocarboximide hydroiodide (3.0 g, 10.4 mmol). The mixture was stirred for 4 hr at 45 °C, dumped into basic water and extracted with ethyl acetate (3 X 100 ml). The combined extracts were washed with water, dried over magnesium sulfate, filtered and concentrated to an oil. The oil was dissolved in methanol, treated with IPA/HCl and triturated with ether. The solids were collected by filtration and washed with ether. One recrystallization from IPA/MeOH/Et₂O yielded a white solid: (2.7 g, 71%), m.p. dec > 260 °C, $[\alpha]_D -44.5^\circ$ (c 0.57 DMSO).

Example 19N-(2,3,4,5-Tetrahydro-1H-3-benzazepine-7-yl)thiophene-2-carboximidamide(a) 2,3,4,5-Tetrahydro-1H-3-benzazepin-7-amine monohydrochloride

To a solution of 2,3,4,5-tetrahydro-7-nitro-1H-3-benzazepine hydrochloride (1.68 g, 7.35 mmol) in ethanol (100 ml) was added 5% palladium on carbon (0.2 g) and the solution was placed on a Paar Hydrogenator Apparatus and pressurized with 45 psi of hydrogen. After the theoretical uptake of hydrogen had been achieved (2 h), the catalyst was filtered off and washed with water (25 ml). The filtrate was concentrated. Absolute ethanol was added and evaporated until all of the water had been evaporated and a solid formed. The solid was dissolved in hot ethanol (50 ml) and the product was precipitated by the addition of ether (75 ml). The solid was collected and air-dried to give the product as an off-white solid (2.43 g (94%)), m.p. 288-91 °C.

(b) N-(2,3,4,5-tetrahydro-1H-3-benzazepine-7-yl)thiophene-2-carboximidamide

-39-

A suspension of 2,3,4,5-tetrahydro-1H-3-benzazepin-7-amine monohydrochloride (0.60 g, 3.0 mmol) and of S-methyl 2-thiophenethiocarboximide hydroiodide (1.1 g, 3.8 mmol) in dimethyl formamide (2.0 ml) and isopropanol (2.0 ml) was stirred at ambience for 20 h. The solid from the reaction was collected and washed with isopropanol (5 ml) and ethyl acetate (15 ml). The air-dried solid weighed 1.18 g and was a mixed salt. This solid was dissolved in water and was basified and extracted into ethyl acetate. The solvent was dried over magnesium sulfate and concentrated to give the free base as a yellow solid. This was taken up in isopropanol (30 ml) and acidified with hydrogen bromide in isopropanol until the solution was acidic. The product was precipitated by the addition of ethyl acetate (35 ml). The product was collected and was dried to give the product as the dihydrobromide salt (0.70 g (54%)), m.p. 281-3 °C.

Example 20

15 N-(1,2,3,4-Tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide

(a) 1,2,3,4-Tetrahydroisoquinolin-7-amine monohydrochloride

This was prepared following the method of Example 19, step (a). From 7-nitro-1,2,3,4-tetrahydroisoquinoline hydrochloride (3.00 g, 14.0 mmol) and 5% palladium on carbon (0.3 g) in ethanol (150 ml) was isolated product as a light rose colored solid (2.43 g (94%)), m.p. 232-4 °C.

(b) N-(1,2,3,4-tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide

This was prepared following the method of Example 19, step (b). From 1,2,3,4-tetrahydroisoquinolin-7-amine monohydrochloride (0.46 g) and S-methyl 2-thiophenethiocarboximide hydroiodide (0.85 g) in isopropanol (2.0 ml) and dimethyl formamide (2.0 ml) was isolated after workup the title compound as the free base (0.60 g (94%)). This was converted to the bisoxalate salt in a methanol/ethyl acetate solution to give the product as an off-white solid (0.59 g (54%)), m.p. 199-200 °C (dec).

30 Example 21

N-(2-benzyl-1,2,3,4-tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide

-40-

(a) 2-benzyl-7-nitro-1,2,3,4-tetrahydroisoquinoline monohydrochloride

To a solution of 7-nitro-1,2,3,4-tetrahydroisoquinoline monohydrochloride (2.50 g, 11.6 mmol) and potassium carbonate (2.0 g) in acetonitrile (100 ml) was added benzyl bromide (2.22 g, 13.0 mmol) in acetonitrile (10 ml). The solution was stirred overnight and the solid was then removed by filtration. The solvent was removed in vacuo to give a solid which was partitioned between methylene chloride and water. The dried (MgSO_4) organic phase was concentrated and the resulting oil was taken up in ethanol (50 ml). This solution was made acidic with hydrochloric acid in ethanol. The precipitate which formed had set up solid and an additional 150 ml of ethanol and 50 ml of ether was added. The solid was collected and air dried to give 2-benzyl-7-nitro-1,2,3,4-tetrahydroisoquinoline hydrochloride as an off white solid (2.78 g (79%)), m.p. 256-8 °C (dec).

(b) 2-benzyl-1,2,3,4-tetrahydroisoquinolin-7-amine hydrochloride

This compound was prepared following the method of Example 19, step (a). From 2-benzyl-7-nitro-1,2,3,4-tetrahydroisoquinoline hydrochloride (2.00 g, 6.56 mmol) and 5% palladium on carbon (0.2 g) in ethanol (100 ml) was isolated the product as a yellow colored solid (1.05 g (78%)), m.p. 257-9 °C (dec).

(c) N-(2-benzyl-1,2,3,4-tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide

This was prepared following the method of Example 19, step (b). From 2-benzyl-1,2,3,4-tetrahydroisoquinolin-7-amine monohydrochloride (0.50 g, 1.8 mmol) and S-methyl 2-thiophenethiocarboximide hydroiodide (0.67 g, 2.3 mmol) in isopropanol (2.0 ml) and dimethyl formamide (2.0 ml) was isolated the title compound as a yellow solid (0.53 g (84%)). This was converted to the oxalate salt in isopropanol, m/e = 348 (M+H).

Example 22

N-(1,2,3,4-tetrahydroisoquinolin-5-yl)thiophene-2-carboximidamide, dioxalate salt

This compound was prepared by following a process analogous to that described above in Example 20. m.p. 75 °C (dec).

Example 23

-41-

N-(1,2,3,4-tetrahydroisoquinoline-6-yl)thiophene-2-carboximidamide(a) 1,2,3,4-tetrahydroisoquinolin-6-amine monohydrochloride

A solution of isoquinolin-6-amine [Manske, R. H. F., et. al., *J. Am. Chem. Soc.*, **72**, 4997 (1950)] (4.40 g, 30.5 mmol) and platinum oxide (300 mg) in a solution of
5 acetic acid (85 ml) and 2.5 M hydrochloric acid (30 ml) was placed on a Paar Hydrogenator Apparatus and was pressurized with 45 psi of hydrogen for 16 h. The solvent was removed in vacuo and the resulting salt was partitioned between aqueous potassium carbonate and 20% isopropanol in methylene chloride. The dried (magnesium sulfate) organic phase was concentrated and the resulting oil was
10 chromatographed on silica gel using 2% methanol in chloroform as eluent to give 3.08 g (93 %) of the product as an oily solid. This product (3.08 g, 20.8 mmol) was taken up in 200 ml of ethanol and 1 equivalent of a 0.1000 M hydrochloric acid solution was added. The solvent was removed to yield the monohydrochloride as a solid, MS 149 (M+H).

15 (b) N-(1,2,3,4-tetrahydroisoquinoline-6-yl)thiophene-2-carboximidamide

This was prepared following the method of Example 19, step (b). From 1,2,3,4-tetrahydroisoquinolin-6-amine monohydrochloride (0.90 g, 4.9 mmol) and S-methyl 2-thiophenethiocarboximide hydroiodide (1.80 g, 6.2 mmol) in isopropanol (2.0 ml) and dimethyl formamide (2.0 ml) was isolated after workup and chromatography on
20 silica gel the title compound as the free base (0.74 g (57%)), m.p. 170-5 °C.

Example 24N-(Isoquinolin-7-yl)thiophene-2-carboximidamide(a) 7-Nitroisoquinoline

25 A solution of 7-nitro-3,4-dihydroisoquinoline (3.00 g, 17.0 mmol) and 5 % palladium on carbon (3.0 g) in decalin (75 ml) was heated at reflux for 3 h. Upon cooling, the solution was filtered and the catalyst washed with chloroform (200 ml). The solvent was removed in vacuo to give 7-nitroisoquinoline (1.63 g) as a tan solid. MS 175 (M+H).

30 (b) Isoquinolin-7-amine

-42-

7-Nitroisoquinoline (1.62 g, 9.25 mmol) in ethanol (150 ml) was hydrogenated on a Paar Hydrogenator Apparatus over 5 % palladium on carbon (0.2 g) as catalyst for 3 h at 50 psi. The reaction mixture was filtered and the solvent removed at reduced pressure. Recrystallization of the solid from ethanol (3 ml) gave
5 isoquinolin-7-amine (0.98 g) as a tan solid. MS 145 (M+H), NMR (CDCl₃) 9.02 (s, 1H), 8.29 (d, 1H), 7.63 (d, 1H), 7.47 (d, 1H), 7.13 (dd, H), 7.03 (d, 1H), 4.00 (broad, 2H).

(c) N-(Isoquinolin-7-yl)thiophene-2-carboximidamide


A solution of isoquinolin-7-amine (0.96 g, 6.7 mmol) and S-methyl 2-
10 thiophenethiocarboximide (2.42 g, 8.36 mmol) in isopropanol (4 ml) and DMF (4 ml) was stirred for 18 h. The solution was poured into dilute sodium hydroxide and extracted with methylene chloride. The extract was dried over magnesium sulfate and the solvent evaporated to give an oil which solidified on standing. The sample
15 was purified by column chromatography on silica gel (5 % methanol in chloroform saturated with gaseous ammonia) to give 1.31 g of a solid. The solid was recrystallized from ethyl acetate (25 ml) to give 1.05 g of the title compound as an off-white solid, m.p. 177.5-8.5 °C.

-43-

Claims

1. A compound of formula I

5



wherein

D represents a five membered heterocyclic aromatic ring containing 1 to 4
10 heteroatoms selected from O, N or S, optionally substituted at a carbon atom by
halogen, trifluoromethyl, alkyl C1 to 6, nitro or cyano, and which is connected to
the remainder of the compound of formula I through a carbon atom;

A represents N(X) or CH(-(CH₂)_m-NXY);

U represents NH, O or CH₂;

15 V represents (CH₂)_a;

W represents (CH₂)_b;

a and b independently represent an integer 0 to 3,

provided that a+b is in the range 1 to 3;

X and Y independently represent hydrogen, alkyl C1 to 6, or the group -(CH₂)_nQ

20 or -NXY represents piperidinyl, pyrrolidinyl, morpholinyl or tetrahydroisoquinolinyl;

Q represents biphenyl or phenyl optionally substituted by one or more groups
selected from alkyl C1 to 6, alkoxy C1 to 6, perfluoroalkyl C1 to 6, halogen, nitro or
cyano;

m represents an integer 0 to 5;

25 n represents an integer 0 to 6;


or the chain U-V-A-W is as defined above save that it may be unsaturated,

or the chain U-V-A-W may represent -NH-CH₂-CH₂-O- substituted at a carbon
atom by the group -(CH₂)_m-NXY, wherein m, X and Y are as defined above,

and pharmaceutically acceptable salts and enantiomers thereof.

30 2. A compound of formula I, according to claim 1, having structure defined by
formula IA:

-44-



5 wherein

T represents a $C_{3,5}$ saturated or unsaturated alkylene chain substituted by $-(CH_2)_m-NXY$; $-O-(CH_2)_2-NH-$ substituted by $-(CH_2)_m-NXY$; or $-U-(CH_2)_a-N(X)-(CH_2)_b-$;

X and Y independently represent hydrogen, alkyl C1 to 6, or the group $-(CH_2)_nQ$, or $-NXY$ represents piperidinyl, pyrrolidinyl, morpholinyl or tetrahydroisoquinolinyl;

10 Q represents phenyl optionally substituted by alkyl C1 to 6, alkoxy C1 to 6, trifluoromethyl, halogen, nitro or cyano;

and U, m, n, a, b and D are as defined above,

save that when T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$ and X represents $-(CH_2)_nQ$, n represents an integer 0 to 5,

15 and pharmaceutically acceptable salts and enantiomers thereof.

3. A compound of formula I according to claim 2, wherein T represents a $C_{3,5}$ saturated or unsaturated alkylene chain substituted by $-(CH_2)_m-NXY$; or $-O-(CH_2)_2-NH-$ substituted by $-(CH_2)_m-NXY$; and X and Y independently represent hydrogen, alkyl C1 to 6 or the group $-(CH_2)_nQ$.

20 4. A compound of formula I, according to claim 2 or claim 3, wherein T represents a $C_{3,5}$ saturated or unsaturated alkylene chain substituted by $-(CH_2)_m-NXY$.

5. A compound of formula I, according to any one of claims 1 to 4, wherein m represents 0 or 1.

25 6. A compound of formula I, according to claim 2, wherein T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$ and X represents hydrogen, alkyl C1 to 6 or the group $-(CH_2)_nQ$.

7. A compound according to claim 2 or claim 6, wherein T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$ and U represents CH_2 .

30 8. A compound according to claim 2, 6 or 7 wherein T represents $-U-(CH_2)_a-N(X)-(CH_2)_b-$ and $a+b$ is 1 or 2.

-45-

9. A compound according to any one of the preceding claims in which n represents 0, 1 or 2 and X and/or Y represents $-(CH_2)_nQ$.
10. A compound according to any one of the preceding claims in which X and/or Y represent $-(CH_2)_nQ$, and Q represents phenyl optionally substituted by alkyl C1 to 6 or halogen.
11. A compound of formula I according to any one of the preceding claims wherein D represents a five membered heterocyclic ring containing one heteroatom selected from O, N or S, optionally substituted at a carbon atom by halogen.
12. A compound of formula I according to claim 11 wherein D represents thienyl, pyrrolyl or furyl.
13. A compound of formula I according to claim 12 wherein D represents 2-thienyl.
14. A compound of formula I which is:
 N-((2-(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide;
 15 N-((2-(phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;
 N-((2-amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;
 N-((1-amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;
 N-((2-amino)indan-5-yl)-2-thiophenecarboximidamide;
 20 N-((2-(methyl)(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide;
 N-((1-amino)indan-6-yl)-2-thiophenecarboximidamide;
 N-((1-(phenylmethyl)amino)indan-6-yl)-2-thiophenecarboximidamide;
 N-((2-((3-chlorophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 N-((2-((2-methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 25 N-((2-((3-methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 N-((2-((4-methylphenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 N-((2-(ethyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 N-((2-(((4-phenyl)phenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 N-((2-(((4-hexyl)phenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;
 30 N-((2-((3-bromophenyl)methyl)amino)indan-5-yl)-2-thiophenecarboxamidine;

-46-

N-((2-((3-chlorophenyl)methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;

N-((2-(phenylmethyl)(methyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;

5 N-((1-(phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-thiophenecarboximidamide;

N-((1-(phenylmethyl)amino)indan-5-yl)-2-thiophenecarboximidamide;

N-((1-(phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-6-yl)-2-thiophenecarboximidamide;

10 N-((2-(phenylmethyl)amino)-1,2,3,4-tetrahydronaphth-7-yl)-2-furancarboximidamide;

N-(2,3,4,5-tetrahydro-1H-3-benzazepine-7-yl)thiophene-2-carboximidamide;

N-(1,2,3,4-tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide;

N-(2-benzyl-1,2,3,4-tetrahydroisoquinoline-7-yl)thiophene-2-carboximidamide;

N-(1,2,3,4-tetrahydroisoquinolin-5-yl)thiophene-2-carboximidamide;

15 N-(1,2,3,4-tetrahydroisoquinoline-6-yl)thiophene-2-carboximidamide;

N-(isoquinolin-7-yl)thiophene-2-carboximidamide;

or a pharmaceutically acceptable salt or enantiomer thereof.

15. A compound according to any one of the preceding claims for use as a pharmaceutical.

20 16. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 14 in admixture with a pharmaceutically acceptable diluent or carrier.

17. The use of a compound as claimed in any one of claims 1 to 14 in the manufacture of a medicament for the treatment or prophylaxis of a disease or
25 condition in which the synthesis or over-synthesis of nitric oxide synthetase forms a contributory part.

18. The use of a compound as claimed in any one of claims 1 to 14 in the manufacture of a medicament for the treatment or prophylaxis of neurodegenerative disorders, of migraine or for the prevention and reversal of
30 tolerance to opiates and diazepines or for the treatment of drug addiction.

-47-


19. A method of treatment or prophylaxis of a disease or condition in which the synthesis or over-synthesis of nitric oxide synthetase forms a contributory part which comprises administering to a person suffering from or susceptible to such a condition a therapeutically effective amount of a compound of formula I as claimed
5 in any one of claims 1 to 14.

20. A method of treatment or prophylaxis of neurodegenerative disorders or of migraine or of prevention and reversal of tolerance to opiates and diazepines or of treatment of drug addiction which comprises administering to a person suffering from or susceptible to such a disease or condition a therapeutically effective amount
10 of a compound of formula I as claimed in any one of claims 1 to 14.

21. A process for the preparation of a compound of formula I, as claimed in claim 1, which comprises:

(a) preparing a compound of formula I by reacting a corresponding compound of formula II


15



I I

wherein U, V, A and W are as defined in claim 1,
with a compound of formula III


20



I I I

wherein D is as defined in claim 1 and L is a leaving group;

(b) preparing a compound of formula I by reacting a corresponding compound
25 of formula IV



I V

30 wherein U, V, A and W are as defined in claim 1 and HA is an acid,
with a compound of formula V




wherein D is as defined in claim 1:

- (c) preparing a compound of formula I in which A represents N(X) and X represents alkyl C1 to 6 or the group $-(CH_2)_nQ$ by reacting a corresponding compound of formula I in which X represents hydrogen with a compound of formula VI




- 10 wherein R⁹ represents alkyl C1 to 6 or the group $-(CH_2)_nQ$ and L is a leaving group;
- (d) preparing a compound of formula I in which A represents $CH(-(CH_2)_m-NXY)$ and at least one of X and Y represents alkyl C1 to 6 or the group $-(CH_2)_nQ$ by reacting a corresponding compound of formula I in which one or both of X and
- 15 Y represents hydrogen with a compound of formula VI;
- (e) preparing a compound of formula I in which A represents $CH(-(CH_2)_m-NXY)$ and m represents an integer 1 to 5, by reduction of a corresponding compound of formula VII



wherein U, V, W, X, Y and D are as defined in claim 1;


- (f) preparing of a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m-$
25 $\text{NXY})$ and both X and Y represent hydrogen, by reduction of a corresponding
compound of formula VIII



wherein U, V, W, m and D are as defined in claim 1;


-49-

(g) preparing a compound of formula I in which A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, X represents hydrogen and m represents an integer 1 to 5, by reduction of a corresponding compound of formula IX




wherein U, V, W, D and Y are as defined in claim 1;

(h) preparing a compound of formula I wherein A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, one of X and Y represents hydrogen, and the other represents $-(\text{CH}_2)_n\text{Q}$ in which n represents an integer 1 to 6, by reduction of a corresponding compound of formula X



wherein Q, m, U, V, W and D are as defined in claim 1;

(i) preparing a compound of formula I wherein A represents $\text{CH}(-(\text{CH}_2)_m-\text{NXY})$, one of X and Y represents hydrogen, and the other represents $-(\text{CH}_2)_n\text{Q}$ in which n represents an integer 1 to 6, by reduction of a compound of formula XI




wherein Q, m, U, V, W and D are as defined in claim 1; or

(j) preparing a compound of formula I in which A represents $\text{CH}(-\text{NXY})$ and X represents hydrogen by reduction of a corresponding compound of formula XII

30

-50-



X I I

- 5 wherein U, V, W, D and Y are as defined in claim 1;
and where desired or necessary converting the resultant compound of formula I, or another salt thereof, to a pharmaceutically acceptable salt thereof, or vice versa.

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/GB 95/01896

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D333/38 A61K31/38 A61K31/47 A61K31/55 C07D307/68
C07D409/12

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,95 05363 (FISONS CORP ; FISONS PLC (GB); GENTILE ROBERT JAMES (US); MURRAY RO) 23 February 1995 cited in the application see abstract; claims; examples 7-9 ---	1-21
A	WO,A,94 21621 (FISONS PLC ; FISONS CORP (US); MACDONALD JAMES EDWIN (US); GENTILE) 29 September 1994 cited in the application see abstract; claims ---	1-21
E	WO,A,96 01817 (ASTRA AB ; MACDONALD JAMES EDWIN (US); SHAKESPEARE WILLIAM CALVIN ()) 25 January 1996 see the whole document -----	1-21

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

4 April 1996

Date of mailing of the international search report

26. 04. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Paisdor, B

INTERNATIONAL SEARCH REPORT

Inter: nal application No.

PCT/GB95/01896

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 19 and 20 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 95/01896

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9505363	23-02-95	AU-B- 7387594 ZA-A- 9406095	14-03-95 19-04-95
WO-A-9421621	29-09-94	AU-B- 6286294 EP-A- 0690851	11-10-94 10-01-96
WO-A-9601817	25-01-96	AU-B- 2413995	09-02-96